



# Investigation of In Vitro Superoxide Scavenging Capacity of Cape

Kadir Batçoğlu\*, I. Çetin Öztürk\*\*

\* Department of Biochemistry, İnönü University Institute of Health Sciences, , Malatya

\*\* Department of Biochemistry, İnönü University School of Medicine, Malatya

**Introduction:** The antioxidant capacity of CAPE (caffeic acid phenethyl ester) has been investigated in recent years but, so far most of the studies were in vivo studies. In this study we investigated, the in vitro antioxidant capacity of CAPE to scavenge superoxide radicals. CAPE, an active component of propolis from honeybee hives, is known to have anti mitogenic, antioxidant, anticarcinogenic, antiinflammatory and immunomodulatory properties.

**Aim of the study:** CAPE is an antioxidant molecule in vivo. The aim of this study is to investigate if CAPE under in vitro conditions has the same effect similar to in vivo.

**Methods:** The method developed by McCord and Fridovich for the determination of SOD activity is modified by replacing SOD with CAPE.

**Results:** We find that, CAPE has superoxide scavenging capacity in a dose dependent manner. The presence of 50 µmol/L CAPE caused 73.66 % percentage of inhibition in cyt c reduction.

**Keywords:** CAPE, Superoxide scavenging capacity, Oxidative stress.

## Cape'nin Süperoksit Süpürücü Kapasitesinin İn Vitro İncelenmesi

**Giriş:** Son yıllarda CAPE (Kafeik asit fenil esterı)'nin antioksidan kapasitesi araştırılmıştır, ancak şu ana kadar yapılmış olan çalışmaların çoğu in vivo çalışmalardır. Biz bu çalışmada CAPE'nin süperoksit radikallerine karşı antioksidan kapasitesini in vitro araştırdık. CAPE bal arısı kovanı propolisinin aktif bir bileşeni olup antimitojenik, antioksidan, antikarsinojen, antiinflamator ve immünomodülatör özellikleri bilinmektedir.

**Amaç:** CAPE in vivo antioksidan bir moleküldür. Bu çalışmada, CAPE'nin benzer koşullarda ancak in vitro ortamda aynı etkiyi gösterip göstermediğinin araştırılması amaçlandı.

**Metod:** McCord ve Fridovich'in SOD aktivitesi tayini için geliştirdikleri yöntem, SOD yerine CAPE kullanılarak modifiye edildi.

**Bulgular:** CAPE'nin doza bağımlı olarak artan bir süperoksit süpürücü kapasiteye sahip olduğunu saptadık. 50 µmol/L CAPE varlığında cyt c redüksiyonu %73,66 inhibe oldu.

**Anahtar Kelimeler:** CAPE, Süperoksit Süpürücü Kapasite, Oksidatif Stres.

Electrons, usually paired one electrically charged and rotate upon themselves while inducing a magnetic field called spin. An electron doublet is more stable than two isolated electrons because the pairing of two electrons with opposite spin cancels their reciprocal magnetic fields.<sup>1</sup>

On the other hand, a free radical is a neutral or charged chemical species whose peripheral shell contains an unpaired electron called, singular electron. Free radicals thus contain an odd number of electrons.<sup>2,3</sup>

Free radicals are produced continuously within the cells. The main free radical generators in cells are the mitochondrial electron transport system, autooxidized molecules such as xanthine oxidase, aldehyde oxidase and

microsomal oxidations.<sup>4</sup> Under conditions such as ischemia/reperfusion, inflammation, xenobiotic metabolism and hyperoxy, excessive reactive oxygen species are produced as a result of oxidative stress leading to damage to all biological molecules.<sup>5-7</sup> The oxygen radicals attack nucleic acids and various modified bases in DNA are generated. Among them, 8-oxoguanine is the most abundant, and appears to play critical roles in carcinogenesis and in aging.<sup>8</sup>

When the delicate balance between the intracellular levels of oxidants and antioxidants is perturbed, ROS (reactive oxygen species) are produced in excess and cells are exposed to oxidative stress.<sup>9</sup> The defence mechanisms of the cells against oxidative stress are superoxide dismutase (SOD), glutathione peroxidase (GPx), and catalase (CAT), antioxidant vitamins such as vitamin A, E and C, antioxidant molecules such as glutathione, melatonin, bilirubin and elements such as Selenium.<sup>10</sup>

Caffeic acid (3,4-dihydroxycinnamic acid) phenethyl ester (CAPE) is an active component of propolis from honeybee hives.<sup>11</sup> It has antiviral, antiinflammatory and immunomodulatory properties and has been shown to inhibit the growth of different types of transformed cells. It has been reported that CAPE suppresses lipid peroxidation and displays antioxidant activity.<sup>12-14</sup> On the other hand Laranjinha et al have reported that CAPE is an important free radical scavenger.<sup>15</sup>

In order to investigate antiinflammatory properties of CAPE, Michaluart et al, made a study on rats and they found that, prostoglandine synthesis was suppressed by high doses of CAPE in acute inflammation.<sup>16</sup> The same researchers reported that, CAPE has an inhibitory effect on cyclooxygenase enzyme. On the other hand, Krol et al studied with ethanolic propolis extracts and they showed that, these extracts had scavenging effects on free radicals.<sup>17</sup>

The aim of this work is to construct an in vitro assay in order to demonstrate the superoxide anion radical scavenging capacity of CAPE. The experiment was planned on the basis of the ability of CAPE to inhibit some observing processes. For this reason, the method developed by McCord and Fridovich for the determination of SOD activity is modified by replacing SOD with CAPE.<sup>18</sup>

## MATERIAL AND METHODS

Xanthine, xanthine oxidase (XOD, from buttermilk), cytochrome c (cyt c, from bovine heart), NaOH,  $\text{KH}_2\text{PO}_4$ ,  $\text{Na}_2\text{HPO}_4$ , EDTA were purchased from Sigma Chemical Company (St. Louise, USA).

Caffeic acid phenethyl ester was synthesized in the Chemistry Laboratory using the method described by Grunberger.<sup>19</sup>

The following solutions were prepared for the assays. And all procedures were done at 37 °C. A Shimadzu 1601 UV/VIS spectrophotometer with a connected PC and a Grand LTD 6G thermostability unit adjusted to  $37 \pm 0.1^\circ\text{C}$  was employed for all spectrophotometric assays.

**Solution A:** 0,76 mg xanthine in 10 ml of 0,001 N. NaOH solution mixed with 24,8 mg cyt c in 100 ml of 50 mM phosphate buffer (pH:7,8) containing 0,1 mM EDTA.

**Solution B:** 0,2 U/ml XOD in 0,1 mM EDTA solution.

**Determination of the rate of cyt c( $\text{Fe}^{+3}$ ) reduction in the presence of CAPE:** 10  $\mu\text{l}$  10% ethanol solution of CAPE (to give a final concentration of 10  $\mu\text{mol/ml}$ ) was added to 750  $\mu\text{l}$  of solution A. After incubation at 37 °C for 30 s. 25  $\mu\text{l}$  of solution B and 15  $\mu\text{l}$  of 50 mM PBS (pH:7,8) were added and the absorbance change at 550 nm was observed for 1 min. The assay was repeated with increased amount of CAPE. Each assay was repeated three times and the arithmetic means were calculated and used. A blank was run by substituting CAPE solution with 10% ethanol solution.

**Determination of uric acid absorbance levels:** Uric acid has an absorbance peak at 292 nm (at pH > 7), (20). 100  $\mu\text{l}$  of reaction mix was added to 900  $\mu\text{l}$  PBS (pH:9,4) at 37 °C and specific absorbance of uric acid was measured at 292 nm.

**Statistical analysis:** Results are given as mean values  $\pm$  standart deviation. Differences between variables were tested for significance by Independent Samples T-Test, using a level of significance of  $p < 0,05$ . Relationship between percentage of inhibition of cyt c reduction and concentration of CAPE was analyzed by Regression Analyze.



**DISCUSSION:**

The role of oxidative stress in many diseases including cancer is gaining importance. A wide range of antioxidants have been proposed for use in the treatment of human diseases.

In recent studies, antioxidant properties of many new molecules have been investigated in order to be used as an antioxidant. One of those molecule is CAPE which is derived from honey bee propolis, attracts attention with its antioxidant property. It has been reported in many studies that CAPE has antiviral, antiinflammatory and immunomodulatory properties in addition to its antioxidant property. Masaaki et al, reported that CAPE has showed anti-tumor promotion activity in their study.<sup>21</sup>

Frenkel et al, studied the antioxidant and anticarcinogenic properties of CAPE.<sup>22</sup> In their study, they exposed Sencar mice to a strong cancer promotion, 12-O-tetradecanoylphorbol-13-acetate and then measured oxidized bases 5-hydroxymethyluracil and 8-hydroxyguanine levels in epidermal tissues of mice. As a result, it was founded that DNA oxidation was blocked by CAPE.

Cengarle et al, researched the antioxidant capacity of propolis extract in poliunsaturated lipidic systems and reported that the antioxidant capacity of propolis extract was better than alpha-tocopherol.<sup>23</sup>

Most of the studies until now performed on CAPE are in vivo and only a few studies exist in vitro.<sup>24-29</sup> Our findings are compatable with the literature. In this study it is aimed to investigate the in vitro effects of CAPE. Our findings clearly exhibited that CAPE inhibited reduction of cyt c by superoxide radicals in a dose- dependent manner. CAPE may possibly exhibited this inhibitor effect by means of two different mechanisms. One of the possible mechanisms is the scavenging effect of CAPE on superoxide radicals produced by xanthine/xanthine oxidase system. The second possible mechanism is inhibitor effect of CAPE on xanthine oxidase enzyme which is responsible in producing superoxide radicals. Thus, superoxide radical production may be decreased.

Xanthine oxidase enzyme produces uric acid together with superoxide radicals. If the amount of uric acid produced by xanthine oxidase is decreased in the presence of CAPE then it may be thought that CAPE has an inhibitor effect on xanthine oxidase. Therefore

the specific absorbance of uric acid was measured at 292 nm. in order to determine whether CAPE has an inhibitor effect on xanthine oxidase or not. And it was observed that in the presence of CAPE there was no significant change at the absorbance levels of uric acid. This result showed that CAPE has no inhibitor effect on xanthine oxidase, so it can be said that CAPE is a strong antioxidant molecule and its antioxidant property comes from its scavenger effect on superoxide radicals.

Antolin et al showed that, melatonin a biomolecule which has free radical scavenging effect like CAPE, increased SOD-mRNA level but, it is unknown that, if CAPE has an effect on mRNA synthesis of antioxidant enzymes like superoxide dismutase, catalase or glutathione peroxidase.<sup>30</sup> Probably, our researches will be focused on that point in the future.

**REFERENCES:**

1. Pierrefiche G., Laborit H., Oxygen free radicals, melatonin and aging. *Experimental Gerontology*, 1995, 30:213-227.
2. Buechter D.D., Free radicals and oxygen toxicity. *Pharm. Res.* 1988, 5:253-260.
3. Halliwell B., Tell me about free radicals, doctor: A review. *J. Royal Soc.Med.* 1989, 82:747-752.
4. Gutteridge J., Halliwell B., Oxygen toxicity, oxygen radicals, transition metals and diseases. Review article. *Biochem. J.* 1984, 219:1-14
5. McCord J.M., Free radicals and myocardial ischemia: Overview and outlook. *Free Rad. Biol Med.* 1988, 4:9-15.
6. Sies H., *Oxidative Stress*, Academic Press Inc. London., 1985, 16-21
7. McCord J.M., Fridovich I., The biology and pathology of oxygen radicals. *Ann. Int. Med.* 1978, 89:122-127.
8. Hayakawa H., Hofer A., Thelander L., Metabolic fate of oxidized guanine ribonucleotides in mammalian cells. *Biochemistry*, 1999, 38:3610-36145
9. Aust S.D., Svingen B.A., Free radical in biological systems. *Acad. Press. London*, 1985, 1-28.
10. Guemouri L., Yues A., Herbert B., Biological variability of SOD, CAT and GPx in blood. *Clin. Chem.* 1991, 37:932-937.
11. Burdock G.A., Review of the biological properties and toxicity of bee propolis. *Food Chem. Toxicol.* 1998, 36:347-363.
12. Kujumgiev A., Tsuetkova I., Bankova V., Antibacterial, antifungal and antiviral activity of propolis of different geographical origin. *J. Ethnopharmacol.* 1999, 64:235-340.
13. Marcucci M.C., Ferreres F., Garcia-Viguera C., Phenolic compounds from Brazilian propolis with pharmacological activities. *J. Ethnopharmacol.* 2001, 74:105-122.
14. Matsushige K, Basnet P, Kadota S., Potent free radical scavenging activity of dicaffeoyl quinic acid derivatives from propolis. *J Trad Med.* 1996, 13:217-228.
15. Laranjinha J., Cadenes E., Redox cycles of CAPE, alpha-tocopherol and ascorbate: Implications for protection of LDL against oxidation. *IUBMB Life*, 1999, 48(1):57-65
16. Michaluart P., Masferrer J.L., Carothers A.M., Subbaramaiah K, Inhibitory effects of CAPE on the activity and expression of cyclooxygenase 2 in human oral epithelial cells and in a rat model of inflammation. *Cancer Res.* 1999,59(10):2347-52.
17. Krol W, Czuba Z, Scheller S, Gabry J., Anti-oxidant property of ethanolic extract of propolis as evaluated by inhibiting the chemiluminescence oxidation of luminol. *Biochem Int.* 1990, 21(4):593-597.
18. McCord J.M., Fridovich I., An enzymatic function for eritrocuprein. *J. Biol. Chem.*, 1969, 244:6049-6055.
19. Grunberger D., Banerjee R., Eisinger K., Oltz EM., Efron L., Preferential cytotoxicity on tumor cells by caffeic acid phenethyl ester isolated from propolis., *Experientia*, 1998, 44:230-232.
20. Lawrence Kaplan, *Clinical Chemistry*, Mosby Press, NY, 1996, pp:501.
21. Nomura M., Kaji A., Wei-ya M., Dong Z., Suppression of cell transformation and induction of apoptosis by caffeic acid phenethyl ester *Molecular Carcinogenesis*, 2001, 31(2):83-89
22. Frenkel K., Wie H., Bhimani R., Ye J., Inhibition of tumor promotor mediated processes in mouse skin and bovine lens by CAPE. *Cancer Res.* 1993, 53(6):1255-61.
23. Cengarle L., Carta A., Tillica G., Antioxidant activity of a Sardinian Propolis. *Riv. Ital. Sostanze Grasse*, 1998, 75:551-557.
24. Suzuki H, Takai M, Koide H., Anti tumor effect of the immunoactivity fraction obtained from Brazilian water soluble pro-polis, given in combination with

## Investigation of In Vitro Superoxide Scavenging Capacity of Cape

- anticancer drug in Ehrlich carcinoma bearing mice. Honeybee Science, 1996, 17:1-6.
25. Basnet P, Matsushige K, Hase K, Kadota S., Four di-O-caffeoyl quinic acid derivatives from propolis. Potent hepatoprotective activity in experimental liver injury models. Biol Pharm Bull. 1996, 19(11):1479-84
  26. Basnet P, Matsushige K, Hase K., Potent antihepatotoxic activity of dicaffeoyl quinic acids from propolis. Biol Pharm Bull. 1996,19(4):655-7
  27. Merino N, Gonzalez R, Gonzalez A, Ramirez D., Histopathological evaluation on the effect of red propolis on liver damage induced by CCl<sub>4</sub> in rats. Arch Med Res. 1996, 27:285-289.
  28. Banskota AH, Tezuka Y, Adnyana IK, Midorikawa K., Cytotoxic, hepatoprotective and free radical scavenging effects of propolis from Brazil, Peru, the Netherlands and China. J Ethnopharmacol. 2000, 72(1-2):239-46.
  29. Maria I, Nieva Moreno, Maria I, Isla, Antonio R., Comparison of the free radical-scavenging activity of propolis from several regions of Argentina. J Ethnopharmacol. 2000, 71(2):109-114
  30. Antolin I, Rodriquez C, Sainz RM, Mayo JC Uria H., Neurohormone melatonin prevents cell damage; effect on gene expression for antioxidant enzymes. FASEB J, 1996, 10:882-890

### Corresponding Address:

Kadir Batçiođlu  
İnönü University, Institute of Health Sciences  
Department of Biochemistry, MALATYA  
E-mail: kbatcioglu@inonu.edu.tr