

5. Tartan Z, Kaşıkçıoğlu H, Yapıcı F, Cam N. Spontaneous coronary artery dissection: a long-term follow-up. *Anadolu Kardiyol Derg* 2007; 7: 436-8.
6. Pierre-Justin G, Pierard LA. Spontaneous coronary artery dissection in an antilles man with acute inferior myocardial infarction. *Int J Cardiol* 2007; 118: 237-40.
7. İyisoy A, Öztürk C, Arslan Z, Çelik T, Ünlü M, Cingöz F, et al. Progressive aortic dissection following RCA instent angioplasty. *Int J Cardiol* 2015; 187: 309-10.
8. İyisoy A, Kurşaklıoğlu H, Köse S, Öztürk C, Amasyalı B, Demirtaş E. Spontaneous intimal dissection in a patient with post-infarct angina: identification with intravascular ultrasound and treatment with coronary stenting. *Jpn Heart J* 2003 ;44:557-64.
9. Karabulut A, Tanriverdi S. Acute coronary syndrome secondary to spontaneous dissection of left internal mammary artery by-pass graft nine years after surgery. *Kardiol Pol* 2011; 69: 970-2.
10. Öztürk C, Çelik T, Demirkol S, Demir M, Balta S, Ünlü M, et al. The healing of spontaneous coronary artery dissection with conservative treatment: When to stop. *Int J Cardiol* 2015; 189: 249-51.
11. Öztürk C, Yıldırım AO, Demir M, Haqmal H, Balta S, Ünlü M, et al. The spontaneous coronary artery dissection may need intervention in the proximal segment of the arteries. *Int J Cardiol* 2016; 202: 943-4.

Address for Correspondence: Dr. Cengiz Öztürk

GATA Kardiyoloji Bölümü, Tefrik Sağlam Cad.

06018 Etilik, Ankara-*Türkiye*

Phone: +90 312 304 42 64 Fax: +90 312 304 42 50

E-mail: drcengizozturk@yahoo.com.tr

©Copyright 2016 by Turkish Society of Cardiology - Available online at www.anatoljcardiol.com

DOI:10.14744/AnatolJCardiol.2016.7120



Author's Reply

To the Editor,

We are pleased by the author's (1) interest in our case report entitled "Recurrent spontaneous dissection affecting different coronary arteries of a young female" published in the February 2016 issue (16: 137-40) of *Anatol J Cardiol*.

The authors proposed that medical treatment may be an option for this case because of spontaneous healing potential of the coronary artery dissection and its recurrent nature. However, it should be accepted that there is no guideline-directed treatment and diagnostic algorithm for spontaneous coronary artery dissection. In large case series, conservative treatment is the preferred strategy for stable patients without ongoing ischemia and if the involved arteries are small or medium sized. Patients with ongoing chest pain, ST elevation, or hemodynamic instability should undergo PCI, particularly when the dissection affects major arteries supplying large areas of the myocardium (2–5). An emergency coronary artery bypass grafting (CABG) should be considered if the dissection extends from the left main into the left anterior descending artery (LAD) and circumflex arteries.

In our case, as shown in the first figure, there is a TIMI 0 flow in LAD after the first septal branch. We first performed PCI to relieve the ongoing ischemia and reduce the infarct size. In the

second episode, the patient suffered acute pulmonary edema treated with initial medical treatment; however, repeat angiogram showed persistent flow-limiting lesion, possibly caused by the intramural hematoma. Because of the life-threatening nature of this condition and hemodynamic instability, we were forced to consider the patient for CABG. In the third episode, the reason behind choosing PCI was the patient's severe ischemia that was unresponsive to medical treatment and compromised hemodynamics, with TIMI I–II flow in the right coronary artery.

Moreover, we accept the role of adjunctive intracoronary imaging, such as optical coherence tomography (OCT) and intravascular ultrasound (IVUS), particularly in diagnosing SCAD subtypes, intramural hematoma, and localizing side branch/true lumen for the intervention (6). However, because of lack of IVUS or OCT facilities in our laboratory at that time, we could not use these techniques.

Necip Ermiş

Department of Cardiology, İnönü University, Turgut Özal Medical Center, Malatya-Turkey

References

1. Ermiş N, Yaşar E, Cansel M. Recurrent spontaneous dissection affecting different coronary arteries of a young female. *Anatol J Cardiol* 2016; 16: 137-8.
2. Saw J. Spontaneous coronary artery dissection. *Can J Cardiol* 2013; 29: 1027-33.
3. Alfonso F, Bastante T, Cuesta J, Rodríguez D, Benedicto A, Rivero F. Spontaneous coronary artery dissection: novel insights on diagnosis and management. *Cardiovasc Diagn Ther* 2015; 5: 133-40.
4. Kansara P, Graham S. Spontaneous coronary artery dissection: case series with extended follow up. *J Invasive Cardiol* 2011; 23: 76-80.
5. Tweet MS, Eleid MF, Best PJ, Lennon RJ, Lerman A, Rihal CS, et al. Spontaneous coronary artery dissection: revascularization versus conservative therapy. *Circ Cardiovasc Interv* 2014; 7: 777-86.
6. Jinnouchi H, Sakakura K, Matsuda J, Wakabayashi Y, Wada H, Momomura S, et al. Recurrent spontaneous coronary artery dissection observed with multiple imaging modalities. *Int Heart J* 2013; 54: 181-3.

Address for Correspondence: Dr. Necip Ermiş

İnönü Üniversitesi Turgut Özal Tıp Merkezi,

Kardiyoloji Bölümü, 44280 Malatya-*Türkiye*

Phone: +90 422 3410660/4508

E-mail: neciperemis@yahoo.com

Effects of cardiopulmonary bypass on new-onset atrial fibrillation

To the Editor,

We read the article titled "SYNTAX score predicts postoperative atrial fibrillation in patients undergoing on-pump isolated coronary artery bypass grafting surgery" that is published in *Anatolian J Cardiol* October 18. Epub ahead of print (1), in which the authors described the effects of SYNTAX score on postoper-