



## PART I: Microwave-Assisted Synthesis of Benzimidazoles: An Overview (Until 2013)

Hasan Küçükbay<sup>1\*</sup>

<sup>1</sup>*Department of Chemistry, Faculty of Arts and Sciences, İnönü University, 44280 Malatya-Turkey.*

**Abstract:** Benzimidazole derivatives are interesting heterocycles and are found in many naturally occurring products and various drugs. They exhibit several pharmacological activities such as antitumor, diuretic, fungicidal, bactericidal, antihelmintic, antiallergic, antihistaminic, vasodilator, hypotensive, spasmolytic activities and local analgesic. Owing to the importance of these heterocyclic compounds, the aim of this review is to present the main aspects of the microwave synthetic methods of the benzimidazole derivatives until 2013.

**Keywords:** Benzimidazoles, benzimidazole synthesis, heterocycles, microwave.

**Cite this:** Küçükbay H. PART I: Microwave-Assisted Synthesis of Benzimidazoles: An Overview (Until 2013). JOTCSA. 2017; 4(1):1-22.

**DOI:** 10.18596/jotcsa.91217.

\*Corresponding author. E-mail: [hkucukbay@inonu.edu.tr](mailto:hkucukbay@inonu.edu.tr). Tel: +90-422-3773881; Fax: +90-422-3410037.

## INTRODUCTION

Benzimidazole is a privileged heterocyclic structure for exploring chemical functionality in biological active molecules (1). They have been studied extensively due to their biological activities such as antiviral against a lot of viruses including HIV (2,3), herpes (HSV-1) (4), RNA (5), influenza (6) and human cytomegalovirus (HCMV) (7), potent antiparasitic agents (8), topoisomerase I inhibitors (9), potential antitumor agents (10), selective neuropeptide Y Y1 receptor antagonists, angiotensin II (AII) inhibitors (11), inhibitors of the hepatitis C virus RNA polymerase (12), antibacterial (13), antifungal (14), antiulcer (15), anti-inflammatory (16), antiprotozoal (17), anticonvulsant (18), antitubercular (19), antidepressant (20), antileishmanial (21), antihypertensive (22), antioxidant (23), and anticoagulant (24, 25).

Moreover, benzimidazole structures behave as good ligands to the transition metal atoms using one of the nitrogen atom and these type of complexes have been found in a variety of biological molecules including ion-heme systems, vitamin B<sub>12</sub> and its derivatives, and several metalloproteins. According to a reported study, metal complexes of biological important ligands are sometimes more effective than the free ligands (26). Therefore, their preparations have been continuously increasing and they have become the most studying heterocyclic motif by synthetic organic chemists and biologists. A number of methods have been developed for the synthesis of compounds containing a benzimidazole moiety (27). The most popular synthetic method of benzimidazoles is the dehydration of 1,2-diaminobenzenes with carboxylic acids under vigorous dehydrating reaction conditions in the presence of strong acids such as polyphosphoric acid, hydrochloric acid, boric acid, or *p*-toluenesulfonic acid (28). However, the use of milder reagents, such as Lewis acids (29), inorganic clays (30), or mineral acids (31), has improved both the yield and purity of this reaction (1a). Another important synthesis method of benzimidazoles is the condensation of 1,2-diaminobenzenes with aldehydes in the presence of an oxidative reagent such as nitrobenzene, benzoquinone, sodium metabisulfite, mercuric oxide, lead tetracetate, iodine, copper (II) acetate, indium perfluorooctane sulfonates, ytterbium perfluorooctane sulfonates, and even air, have been used for this aim (32). The benzimidazoles could also be synthesized by the coupling of 1,2-diaminobenzenes with carboxylic acid derivatives such as anhydrides, lactones, orthoesters, imidates and nitriles (33). Moreover, several other catalysts, namely iodine (34), hydrogen peroxide (35), boron trifluoride diethyl etherate (36), zirconyl (IV) chloride (37), zeolite (38), and L-proline (39) have been effectively used for the synthesis of benzimidazole derivatives.


Despite their high effectiveness for the synthesis of benzimidazoles, some of the methods need to be improved for the high reaction temperature, long reaction times, toxic solvent and high-priced catalyst, etc. (40). Therefore, development of mild, efficient and environmentally benign protocol for the synthesis of benzimidazoles is still a hot topic for researchers. After the first reports of applications of microwaves in synthetic chemistry in 1986 (41), microwave-assisted

synthesis have become popular, particularly, last two decades due to the generally short reaction times, the high purity and yields of the resulting products. Up to now, several microwave-assisted methodologies for the synthesis of benzimidazoles have been reported.

This article aims to review the work reported, microwave-assisted synthesis of benzimidazole derivatives during past years until 2013.


### Microwave-assisted benzimidazole synthesis: a) Benzimidazole synthesis from carboxylic acids/carboxylic acid derivatives and 1,2-diaminobenzenes

The first microwave-assisted benzimidazole synthesis reported in 1995 by Bougrin and Soufaoui (42). They have described a benzimidazole synthesis method using 1,2-diaminobenzene or 4-substituted-1,2-diaminobenzene and ethyl acetoacetate or ethyl benzoylacetate on solid mineral supports in dry media under microwave irradiation in domestic ovens (Scheme 1).




**Scheme 1.**

Bougrin *et al.* (43) have described in dry media synthesis of 2-trifluoromethylbenzimidazoles through cyclocondensation of N-(carbotrifluoromethyl)-*ortho*-arylenediamines on montmorillonite K10 under domestic microwave oven with good yields (Scheme 2).




**Scheme 2.**

Bendale and Sun (44) have described an efficient, facile, and practical liquid-phase combinatorial synthesis of benzimidazoles under microwave irradiation in short time (Scheme 3).




**Scheme 3.**

Getvoldsen *et al.* (45) developed an *in situ* monitoring method for the reactions in a microwave using UV/Vis spectroscopy. The benzimidazole synthetic reaction from 1,2-diaminobenzene and formic acid has chosen as a model reaction. They have reported that the new method would be an excellent analytical tool for monitoring the progress of a reaction, determination of end points and derivation of quantitative and kinetic data (Scheme 4).




**Scheme 4.**

Boufatah *et al.* (46) have reported the preparation of some new biologically active benzimidazole-4,7-dione derivatives in 7 steps through the microwave irradiation for the ring closing step to obtain benzimidazole derivative (Scheme 5).




Scheme 5.

Zhang and Tempest (47) have improved the efficiency of the Ugi/de-Boc/cyclization synthesis replaced the normal Boc group with the fluorous-Boc group to protect the diamine. They were used fluorous component as a limiting agent in the Ugi reaction. Hence, they have easily separated designed condensation product containing the F-Boc group from the reaction mixture by fluorous SPE. Compared to the original Ugi/de-Boc/cyclization procedures, which take 1-2 days, the fluorous/microwave approach which modified by Zhang and Tempest has more favorable reaction and purification conditions: less than 20 min for each reaction and no need of the double scavenging step (Scheme 6).




Scheme 6.

Getvoldsen *et al.* (48) have reported 2-([4-<sup>18</sup>F] fluorophenyl) benzimidazole synthesis from the cyclocondensation reaction of 1,2-diaminobenzene with radiolabelled [4-<sup>18</sup>F]fluorobenzoic acid in neat methanesulfonic and polyphosphoric acids under microwave (Scheme 7).




Scheme 7.

Martinez-Palou *et al.* (49) have described 2-long alkyl chain substituted benzimidazole synthesis from the reaction of 1,2-diaminobenzene and stearic acid via mono and multimode microwave irradiation with high yields (Scheme 8).




**Scheme 8.**

Vliet *et al.* (50) have described a simple, rapid, one pot procedure for the generation of 2-substituted benzimidazoles with high-yield directly from 2-nitroanilines using  $\text{SnCl}_2$  as a reduction agent and carboxylic acid under microwave irradiation at 130 °C in 5 min (Scheme 9).




**Scheme 9.**

Lin *et al.* (51) have reported a microwave-assisted one-pot synthesis of several benzimidazole derivatives from readily available starting compounds such as 1,2-diaminobenzene, 4,5-diaminopyrimidine, *cis*-1,2-diaminocyclohexane and several carboxylic acids including heteroaraomatic carboxylic acids (Scheme 10).



**Scheme 10.**


Mobinikhaledi and co-workers (38) described a selective synthesis of 2-aryl-1*H*-benzimidazoles through the reaction of 1,2-diaminobenzene with arylcarboxylic acid using zeolite as an efficient catalyst under microwave irradiation (Scheme 11).



Ar= 3-Nitrophenyl, 4-nitrophenyl, 4-chlorophenyl  
 3-chlorophenyl, 4-tolyl, 3-tolyl, 4-methoxyphenyl  
 4-bromophenyl, 2-pyridyl


**Scheme 11.**

Jing *et al.* (52) have described a simple and rapid synthesis of benzimidazoles from 1,2-diaminobenzene dihydrochloride and esters under microwave irradiation. This is also the first report for microwave assisted benzimidazole synthesis from esters and 1,2-diaminobenzenes (Scheme 12).




**Scheme 12.**

Gellis *et al.* (53) have reported the synthesis a series of variously substituted benzimidazole-4,7-diones in seven steps and antitumor activity results of them. One of them exhibit excellent cytotoxicity comparable to that of mitomycin C (Scheme 13).




**Scheme 13.** Reagents and conditions: (a) KOH,  $(\text{CH}_3)_2\text{SO}_4$ ,  $\text{CH}_3\text{OH}$ , reflux, 1 h; (b) R= H;  $\text{HNO}_3$  62%, 1 h at rt, 1 h at  $100^\circ\text{C}$  or R=  $\text{CH}_3$ ,  $(\text{CH}_3\text{CO})_2\text{O}$ ,  $\text{HNO}_3$  fuming,  $90^\circ\text{C}$ , 1 h; (c) Sn, HCl; (d)  $\text{HOCH}_2\text{CO}_2\text{H}$ , 4 N HCl, microwave, 800 W, 1 h 30 min; (e)  $\text{LiN}(\text{TMS})_2$ , THF, 20 min, then  $\text{CH}_3\text{I}$ , 7h; (f)  $\text{SOCl}_2$ ,  $\text{CHCl}_3$ , reflux, 4h; (g)  $\text{BBr}_3$ ,  $\text{CH}_2\text{Cl}_2$ ,  $-78^\circ\text{C}$ , 3 h; (h)  $\text{K}_2\text{Cr}_2\text{O}_7$ ,  $\text{H}_2\text{O}$ , rt, 2 h.

Algül *et al.* (54) have described the synthesis of some 2-substituted benzimidazole, benzothiazole, and indole derivatives using on both microwave irradiation and conventional heating methods. They have also evaluated hyaluronidase inhibitory activity of the compounds (Scheme 14).




**Scheme 14.** A= PPA/MW, D=  $\text{Na}_2\text{S}_2\text{O}_5/\text{DMF}$ , E= PPA, G= HCl.

Keri *et al.* (27) have reported an expeditious, novel, and efficient method for the synthesis of benzimidazole derivatives from *o*-phenylenediamines with different phenoxy acid using a Wells-Dawson  $\text{H}_6\text{P}_2\text{W}_{18}\text{O}_{62} \cdot 24 \text{H}_2\text{O}$  catalyst (Scheme 15).




Scheme 15.

Similar to the Keri method, Hosamani and co-workers (55) described a convenient protocol for the preparation of 5-nitro-2-aryloxy-1H-benzimidazole libraries in one pot-synthesis both under microwave irradiation and conventional heating methods using hydrochloric acid as catalyst (Scheme 16).




Scheme 16.

Hasaninejad *et al.* (56) have reported the synthesis of some 2-substituted benzimidazole derivatives from benzene-1,2-diamine with mono and dicarboxylic acids under microwave irradiation using silphox [POCl<sub>3-n</sub>(SiO<sub>2</sub>)<sub>n</sub>] catalyst in highly yield and short reaction times (Scheme 17).




Scheme 17.

Sun and co-workers (57) synthesized angular bis-benzimidazoles on the support with three appendages with good yields and purities through soluble polymer-supported reaction. The integration of microwave irradiation and a soluble polymer support strategy provided an efficient and convenient approach for high throughput and diversity-oriented synthesis of drug-like molecules. Their preliminary screening results have shown that some of these compounds exhibited moderately to good inhibition against VEGFR 3, which is related to the invasion and migration of cancer cells (Scheme 18).




Scheme 18.

Sun and co-workers (58-60) reported a novel synthetic method for the substituted benzimidazole derivatives from the starting compounds 1,2-diaminobenzene derivative on soluble polymer support and 3-nitro-4-hydroxy/fluorobenzoic acid in a multistep process under focused microwave irradiation (Scheme 19).


**Scheme 19.**

The same group has recently described the synthesis of bis-heterocyclic skeletal novel benzimidazole-linked pyrrolo-/pyridobenzimidazolones and benzimidazole-linked isoindolo-benzimidazoleones on ionic liquid support under microwave irradiation by utilizing the cascade cyclization (61) (Scheme 20).

**Scheme 20.**

### b) Benzimidazole synthesis from aldehydes and 1,2-diaminobenzenes


Ben-Alloum *et al.* (62) have described oxidative heterocyclization of aldehydes and *o*-phenylenediamine with nitrobenzene or dimethylsulfoxide impregnated on silica gel irradiated with microwave in good yields and high purity (Scheme 21).



R= Ph, 4-tolyl, 4-anisoyl, 4-ClPh, 4-NO<sub>2</sub>Ph, 2-ClPh, 2-HOPh,  
2-furyl, *n*-propyl, ethyl


**Scheme 21.**

Navarrete-Vazquez and co-workers (63) described a simple, rapid, and efficient method for the preparation of several 2-(alkoxyaryl)-1*H*-benzimidazole compounds under solvent-free conditions using readily available and inexpensive reagents utilizing microwave irradiation (Scheme 22).



**Scheme 22.**

Zahran and co-workers (64) described the synthesis of new heterocyclic compounds containing pyrazol-5-one coupled with benzimidazole under dry media and they also explored antitumor activity of the newly synthesized heterocycles. Some of them were found to be more effective than thalidomide (Scheme 23).



**Scheme 23.**


Dubey and Moorthy (65) have reported comparative synthesis of 2-alkyl and 2-aryl substituted benzimidazole derivative in the presence of polyphosphoric acid through microwave and conventional methods and also studied the effect of salt form of reactant for completion of the reaction. According to their report, the 2-substituted aryl and alkyl benzimidazole derivative synthesized via microwave and it is beneficial, in respect of yield increases up to 10 to 50% and time (96 to 98% was reduced) than conventional method of synthesis. They also showed some remarkable inferences while the use of salt form of the reactant (1,2-diaminobenzene dihydrochloride). Navarrete-Vazquez *et al.* (66) have synthesized several benzimidazole derivatives which could be vasorelaxant and spasmolytic properties from the reaction of 1,2-

diaminobenzene with aromatic substituted aldehydes and sodium metabisulfite under microwave irradiation.

Jacob and co-workers (67) described an improved methodology for the selective synthesis of 1,2-disubstituted benzimidazoles by the condensation of 1,2-diaminobenzene and aldehydes using solid-supported catalyst ( $\text{SiO}_2/\text{ZnCl}_2$ ) in a short time under microwave irradiation.


Mao *et al.* (68) have reported intermittent microwave irradiation for the synthesis 2-benzimidazole derivatives from aromatic aldehydes and 1,2-diaminobenzene only using molecular oxygen as the oxidant and inexpensive KI as the catalyst.

Rathod and co-workers (40) described eco-friendly benzimidazole synthesis from aromatic aldehydes and 1,2-diaminobenzene using  $\text{MoO}_3/\text{CeO}_2.\text{ZrO}_2$  as a catalyst under solvent-free conditions both conventional and microwave heating. Azarifar *et al.* (69) have reported 2-aryl-1-(arylmethyl)1H-benzimidazole synthesis from 1,2-diaminobenzene and aldehydes through acetic acid-promoted condensation in air under microwave irradiation (Scheme 24).



**Scheme 24.**


Mungra and co-workers (70) described a microwave-assisted synthesis of 4-(benzimidazol-2-yl)tetrazolo[1,5-a]quinoline from 1,2-diaminobenzene and heterocyclic aldehydes catalyzed by an organocatalyst *p*-TsOH (Scheme 25).



**Scheme 25.**

Krim *et al.* (71) have reported efficient synthetic routes to prepare fluorescent compounds containing benzimidazoles in the 5-position of pyrimidine nucleosides. Their synthesis is based on the straightforward condensation of 5-formyl-2'-deoxyuridine and arylenediamine


derivatives under a cooperative effect of microwave activation and NaHSO<sub>3</sub> catalysis (Scheme 26).



**Scheme 26.**


### c) Miscellaneous benzimidazole syntheses

Reddy *et al.* (72) have reported microwave assisted synthesis of 5- or 5,6-disubstituted-2-trifluoroacetylbenzimidazole derivatives (Scheme 27).




**Scheme 27.**

Hulme and co-workers (73) described a novel two-step solution phase protocol for the synthesis of arrays of triazadibenzoazulenones using the Ugi reaction and two tandem ring closing transformations. They also used microwave irradiation for 5 or 10 min for the second ring closing steps (Scheme 28).



**Scheme 28.**

Wang *et al.* (74) reported microwave-assisted solvent-dependent chemoselective reaction dealing with 4-arylidene-2-phenyloxazol-5-ones and *ortho*-diamines to obtain benzimidazole scaffold (Scheme 29).




R= H, Me, Cl,  $\text{CO}_2\text{H}$

Ar= Ph, 4-ClPh, 4-MeOPh, 3,4,5-trimethoxyphenyl, 2-thienyl

**Scheme 29.**

Shaw and *et al.* (75) described a novel synthetic protocol for the synthesis of unique norstatine analogs bearing benzimidazole moiety through Passerini reaction and following ring closing (Scheme 30).




R<sup>1</sup>=Me, isopropyl, benzyl

R<sup>2</sup>= *n*-propyl, Ph, 4-BrPh, 4-MeOPh, 1-naphthyl

**Scheme 30.**

Wen *et al.* (76) have reported on eco-friendly, one-pot efficient propylphosphonic anhydride (T3P) mediated synthesis of benzimidazoles from various carboxylic acids and 1,2-diaminobenzene under microwave irradiation (Scheme 31).

**Scheme 31.****CONCLUSION**

Properties of benzimidazole and its derivatives have been studied for over a hundred years. Special interest of researchers triggered by the fact that 5,6-dimethylbenzimidazole is a component of naturally occurring vitamin B-12. The benzimidazole ring is an important pharmacophore in modern drug discovery. On the other hand, in the past decade, heating and driving chemical reaction by microwave energy has been an increasingly popular theme in the scientific community. A number of methods have been developed for the synthesis of benzimidazole containing compounds. There are two main synthesis methods for the benzimidazoles: i) from the reaction of 1,2-diaminobenzenes with carboxylic acids or its derivatives, ii) condensation of 1,2-diaminobenzenes with aldehydes presence of an oxidative reagent. Despite their high efficiency for the synthesis of benzimidazoles, some of the methods are plugged by one of the drawbacks such as, high reaction temperature, prolonged reaction times, toxic solvent and expensive catalyst, etc. Therefore, development of mild, efficient and environmentally benign protocol for the synthesis of benzimidazoles continues to attract researchers' attentions. This article aims to review the work reported, microwave-assisted synthesis of benzimidazole derivatives published in SCI journals from the first report up to 2013.

**ABBREVIATIONS**

Ac= Acetyl  
 Ar= Aryl  
 Boc= *tert*-Butyloxycarbonyl  
 DCC= N,N'-Dicyclohexylcarbodiimide  
 DCE= 1,2-Dichloroethane  
 DCM= Dichloromethane  
 DEAD= Diethyl azodicarboxylate  
 DIPEA= N,N-Diisopropylethylamine  
 DMAD= Dimethylacetylene dicarboxylate  
 DMAP= 4-Dimethylaminopyridine  
 DMF= Dimethylformamide  
 DMSO= Dimethyl sulfoxide  
 HCMV= Human cytomegalovirus  
 HIV= Human Immunodeficiency Virus  
 HSV= Herpes simplex virus  
 Me= Methyl  
 MW= Microwave

PEG= Polyethylene glycol  
 PPA= Polyphthalamide  
 RNA= Ribonucleic acid  
 TBDMS= *tert*-butyldimethylsilyl  
 TFA= Trifluoroacetic acid  
 TEA= Triethanolamine  
 Ts= Tosyl  
 T3P= propylphosphonic anhydride  
 VEGFR= Vascular endothelial growth factor receptor

## REFERENCES

1. (a) Zhang Z-H, Yin L, Wang Y-M. An expeditious synthesis of benzimidazole derivatives catalyzed by Lewis acids. *Catalysis Communications*. 2007 Jul;8(7):1126–31. DOI:10.1016/j.catcom.2006.10.022. 1. (b) S Panda S, Malik R, C Jain S. Synthetic approaches to 2-arylbenzimidazoles: A review. *Current Organic Chemistry*. 012;16(16):1905–19.
2. Porcari AR, Devivar RV, Kucera LS, Drach JC, Townsend LB. Design, synthesis, and antiviral evaluations of 1-(substituted benzyl)-2-substituted-5, 6-dichlorobenzimidazoles as nonnucleoside analogues of 2, 5, 6-trichloro-1-( $\beta$ -D-ribofuranosyl) benzimidazole. *Journal of medicinal chemistry*. 1998;41(8):1252–62. DOI: 10.1021/jm970559i.
3. Roth M, Morningstar ML, Boyer PL, Hughes SH, Buckheit Jr RW, Michejda CJ. Synthesis and biological activity of novel nonnucleoside inhibitors of HIV-1 revers transcriptase. 2-aryl-substituted benzimidazoles. *Journal of Medicinal Chemistry* 1997; Dec; 40(26), 4199-4207. DOI:10.1021/jm970096g.
4. Migawa MT, Girardet J-L, Walker JA, Koszalka GW, Chamberlain SD, Drach JC, et al. Design, synthesis, and antiviral activity of  $\alpha$ -nucleosides: D-and L-isomers of Iyxofuranosyl-and (5-deoxylyxofuranosyl) benzimidazoles. *Journal of Medicinal Chemistry*. 1998;41(8):1242–51. DOI: 10.1021/jm970545c.
5. Tamm, I.; Sehgal, P.B. Halobenzimidazole ribosides and RNA synthesis of cells and viruses. *Advances in Virus Research*. 1978, 22, 187-258. DOI: 10.1016/S0065-3527(08)60775-7.
6. Tamm, I. Ribonucleic acid synthesis and influenza virus multiplication. *Science*, 195; 126: 1235-1236.
7. Kim JS, Gatto B, Yu C, Liu A, Liu LF, La Voie EJ. Substituted 2,5'-bi-1H-benzimidazoles: Topoisomerase I inhibition and cytotoxicity. *Journal of Medicinal Chemistry*. 1996; 39: 992-998. DOI: 10.1021/jm950412w.
8. Zarrinmayeh H, Nunes AM, Ornstein PL, Zimmerman DM, Arnold B, Schober DA, Gackenheimer SL, Bruns RF, Hipskind PA, Britton TC, Cantrell BE, Gehlert DR. Synthesis and evaluation of a series of novel 2-[(4-chlorophenoxy)methyl]benzimidazoles as selective neuropeptide Y Y1 receptor antagonists. *Journal of Medicinal Chemistry*. 1998; 41: 2709-2719. DOI: 10.1021/jm9706630.
9. Kohara Y, Kubo K, Imamiya E, Wada T, Inada Y, Naka T. Synthesis and angiotensin II receptor antagonistic activities of benzimidazole derivatives bearing acidic heterocycles as novel tetrazole bioisosteres. *Journal of Medicinal Chemistry*.1996; 39: 5228-5235. DOI: 10.1021/jm960547h.
10. a) Denny WA, Rewcastle GW, Baguley BC. Potential antitumor agents .59. structure-activity-relationships for 2-phenylbenzimidazole-4-carboxamides, a new class of minimal DNA-intercalating agents which may not act via topoisomerase-II. *Journal of Medicinal Chemistry*. 1990; 33: 814-819. DOI: 10.1021/jm00164a054. b) Demirayak S, Mohsen UA, Karaburun AC. Synthesis and anticancer and anti-HIV testing of some pyrazino[1,2-a]benzimidazole derivatives. *European Journal of Medicinal Chemistry*. 2002; 37: 255-260. DOI: 10.1016/S0223-5234(01)01313-7. c) Sharma PS, Sharma R, Tyagi R. Inhibitors of cyclin dependent kinases: Useful targets for cancer treatment. *Current Cancer Drug Targets*. 2008; 8: 53-75. DOI: 10.2174/156800908783497131.
11. Fonseca T, Gigante B, Gilchrist TL. A short synthesis of phenanthro[2,3-d]imidazoles from dehydroabietic acid. Application of the methodology as a convenient route to benzimidazoles. *Tetrahedron*. 2001; 57: 1793-1799. DOI: 10.1016/S0040-4020(00)01158-3.

12. LaPlante SR, Jakalian A, Aubry N, Bousquet Y, Ferland J-M, Gillard J, Lefebvre S, Poirer M, Tsantrizos YS, Kukolj G, Beaulieu PL. Binding mode determination of benzimidazole inhibitors of the hepatitis C virus RNA polymerase by a structure and dynamics strategy. *Angewandte Chemie International Edition*. 2004; 43: 4306-4311. DOI: 10.1002/anie.200460326.
13. a) Fekner T, Gallucci J, Chan MK. Ruffling-induced chirality: Synthesis, metalation, and optical resolution of highly nonplanar, cyclic, benzimidazole-based ligands. *Journal of American Chemical Society*. 2004; 126: 223-236. DOI: 10.1021/ja030196d. b) Küçükbay H, Çetinkaya E, Durmaz R. Synthesis and antimicrobial activity of substituted benzimidazole, benzothiazole and imidazole derivatives. *Arzneimittel Forschung/Drug Research*. 1995; 45: 1331-1334. c) Küçükbay H, Durmaz R, Orhan E, Günal S. Synthesis, antibacterial and antifungal activities of electron-rich olefins derived benzimidazole compounds. *Il Farmaco*. 2003; 58: 431-437. DOI: 10.1016/S0014-827X(03)00068-5 d) Durmaz R, Köroğlu M, Küçükbay H, Temel İ, Özer MK, Refiq M, Çetinkaya E, Çetinkaya B, Yoloğlu S. Investigation of serum minimal inhibitory concentrations of some benzimidazole, imidazole and benzothiazole derivatives and their effects on liver and renal functions. *Arzneimittel Forschung/Drug Research*. 1998; 48: 1179-1184.
14. a) Olander D, Zwawiak J, Lukianchuk V, Lesyk R, Kropacz A, Fojutowski A, Zaprutko L. Synthesis of some N-substituted nitroimidazole derivatives as potential antioxidant and antifungal agents. *European Journal of Medicinal Chemistry*. 2009; 44: 645-652. b) Küçükbay H, Durmaz B. Antifungal activity of organic and organometallic derivatives of benzimidazole and benzothiazole. *Arzneimittel Forschung/Drug Research*. 1997; 47: 667-670.
15. a) Carcangue D, Shue YK, Wuonola MA, Nickelsen MU, Joubran C, Abedi JK, Jones J, Kuhler TC. Novel structure's derived from 2-[[[(2-Pyridyl)methyl]thio]- 1H-benzimidazole as anti-Helicobacter pylori agents, part 2. *Journal of Medicinal Chemistry*. 2002; 45: 4300-4309. DOI: 10.1021/jm020868v. b) Lindberg P, Nordberg P, Alminger T, Bradstrom A, Wallmark B. The mechanism of action of the gastric-acid secretion inhibitor omeprazole. *Journal of Medicinal Chemistry*. 1986; 29: 1327-1329. DOI: 10.1021/jm00158a001.
16. a) Medzhitov R. Inflammation 2010: New adventures of an old flame. *Cell*; 2010;140: 771-776. DOI: 10.1016/j.cell.2010.03.006. b) Grivennikov S, Greten FR, Karin M. Immunity, inflammation, and Cancer. *Cell*. 2010; 140:883889. DOI: 10.1016/j.cell.2010.01.025. c) Nathan C. Points of control in inflammation. *Nature*. 2002; 420: 846-852. DOI: 10.1038/nature01320. d) Cronstein BN, Weissmann G. Targets for antiinflammatory drugs. *Annual Review of Pharmacology and Toxicology*. 1995; 35: 449-462. DOI: 10.1146/annurev.pa.35.040195.002313.
17. a) Vazquez GN, Vilehis MMR, Mulia LY, Melendez V, Gerena L, Campos AH, Castillo R, Luis FH. Synthesis and antiprotozoal activity of some 2- (trifluoromethyl)-1H-benzimidazole bioisosteres. *European Journal of Medicinal Chemistry*. 2006; 41: 135-141. DOI:10.1016/j.ejmech.2005.09.001. b) Gomez HT, Nunez EH, Rivera IL, Alvarez JG, Rivera RC, Puc RM, Ramos RA, Guttirez MCR, Bacab MJC, Vazquez GN. Design, synthesis and in vitro antiprotozoal activity of benzimidazole-pentamidine hybrids. *Bioorganic and Medicinal Chemistry Letters*, 2008; 18: 3147-3151. DOI:10.1016/j.bmcl.2008.05.009.
18. Singh J, Grover P, Pathak DP. Synthesis, anticonvulsant activity and comparative QSAR study of some novel 1,2,5-trisubstituted benzimidazole derivatives. *Acta Pharmaceutica Scientia*. 2010; 52: 511-522.
19. a) Gupta P, Hameed S, Jain R. Ring-substituted imidazoles as a new class of anti-tuberculosis agents. *European Journal of Medicinal Chemistry*. 2004; 39: 805-814. DOI: 10.1016/j.ejmech.2004.05.005. b) Shingalapur RV, Hosamani KM, Keri RS. Synthesis and evaluation of in vitro anti-microbial and anti-tubercular activity of 2-styryl benzimidazoles. *European Journal of Medicinal Chemistry*. 2009; 44: 4244-4248. DOI: 10.1016/j.ejmech.2009.05.021. c) Jyoti P, Vinod TK, Shyam VS, Vinita C, Bhatnagar S, Sinha S, Gaikwad AN, Tripathi RP. Synthesis and antitubercular screening of imidazole derivatives. *European Journal of Medicinal Chemistry*. 2009; 44: 3350-3355. DOI: 10.1016/j.ejmech.2009.02.013.
20. Hadizadeh F, Hosseinzadeh H, Sadat M-SV, Seifi M, Kazemi S. Synthesis and antidepressant activity of N-substituted imidazole-5-carboxamides in forced swimming test model. *Iranian Journal of Pharmaceutical Research*. 2008; 7: 29-33.
21. Bhandari K, Srinivas N, Marrapu VK, Verma A, Srivastava S, Gupta S. Synthesis of substituted aryloxy alkyl and aryloxy aryl alkyl imidazoles as antileishmanial agents. *Bioorganic and Medicinal Chemistry Letters*. 2010; 20: 291-293. DOI:10.1016/j.bmcl.2009.10.117.

22. a) Naik P, Murumkar P, Giridhar R, Yadav MR. Angiotensin II receptor type 1 (AT(1)) selective nonpeptidic antagonists-A perspective. *Bioorganic and Medicinal Chemistry*. 2010; 18: 8418-8456. DOI: 10.1016/j.bmc.2010.10.043. b) Kubo K, Inada Y, Kohara Y, Sugiura Y, Ojima M, Itoh K, Furukawa Y, Nishikawa K, Nakat T. Nonpeptide angiotensin-II receptor antagonists-synthesis and biological-activity of benzimidazoles. *Journal of Medicinal Chemistry*. 1993; 36: 1772-1784. DOI: 10.1021/jm00064a011.
- 23 a) Cole ER, Crank G, Salam-Sheikh A. Antioxidant properties of benzimidazoles. *Journal of Agricultural and Food Chemistry*. 1974; 22: 918. b) Neochoritis CG, Zarganes-Tzitzikas T, Tsoleridis CA, Stephanidou- Stephanatou J, Kontogiorgis CA, Hadjipavlou-Litina DJ, Choli-Papadopoulou T. One-pot microwave assisted synthesis under green chemistry conditions, antioxidant screening, and cytotoxicity assessments of benzimidazole Schiff bases and pyrimido[1,2-a]benzimidazol-3(4H)-ones. *European Journal of Medicinal Chemistry*. 2011; 46: 297-306. DOI: 10.1016/j.ejmech.2010.11.018.
24. Huel NH, Nar H, Priepke H, Ries U, Stassen JM, Wiene W. Structure-based design of novel potent nonpeptide thrombin inhibitors. *Journal of Medicinal Chemistry*. 2002; 45: 1757-1766. DOI: 10.1021/jm0109513.
25. Evans BE, Rittle KE, Bock MG, Dipardo RM, Freidinger RM, Whitter WL, Lundell F, Veber DF, Anderson PS, Chang RS, Chang RSL, Lotti VJ, Gerino DJ, Chen TB, Kling PJ, Kunkel KA, Springer JP, Hirshfield J. Methods for drug discovery: development of potent, selective, orally effective cholecystokinin antagonists. *Journal of Medicinal Chemistry*. 1998; 42: 2235-2246. DOI: 10.1021/jm00120a002.
26. Ahuja IS, Prasad I. Isonicotinamide complexes with some metal(II) halides and pseudohalides, *Inorganic and Nuclear Chemistry Letters*. 1976; 12: 777-784. DOI:10.1016/0020-1650(76)80065-7.
27. Keri RS, Hosamani KM, Seetharama HR, Reddy RS, Shingalapur RV. Wells-Dawson heteropolyacid: An efficient recyclable catalyst for the synthesis of benzimidazoles under microwave condition. *Catalyst Letter*. 2009; 131: 552-559. DOI: 10.1007/s10562-009-9966-2.
28. Jing X, Zhu Q, Xu F, Ren X, Li D, Yan C. Rapid one-pot preparation of 2-substituted benzimidazoles from esters using microwave conditions. *Synthetic Communication*. 2006; 36: 2597-2601.
29. Tandon VK, Kumar M. BF<sub>3</sub> center dot Et<sub>2</sub>O promoted one-pot expeditious and convenient synthesis of 2-substituted benzimidazoles and 3,1,5-benzoxadiazepines. *Tetrahedron Letters*. 2004; 45: 4185-4187.
30. Bougrin K, Loupy A, Petit A, Daou B, Soufiaoui M. Novel synthesis of 2-trifluoromethylarylimidazoles on montmorillonite K10 in a 'dry medium' under microwave irradiation. *Tetrahedron*. 2001; 57: 163-168.
31. Rastogi R, Sharma S. 2-Aminobenzimidazoles in Organic Synthesis. *Synthesis*. 1983; 861-882. DOI: 10.1055/s-1983-30546.
32. Hegedus A, Hell Z, Potor A. Zeolite-catalyzed environmentally friendly synthesis of benzimidazole derivatives. *Synthetic Communications*. 2006; 36: 3625-3630. DOI:10.1080/00397910600943865.
33. Lin SY, Isome Y, Stewart E, Liu JF, Yohannes D, Yu L. Microwave-assisted one step high-throughput synthesis of benzimidazoles. *Tetrahedron Letters*. 2006; 47: 2883-2886. DOI: 10.1016/j.tetlet.2006.02.127.
34. Sun P, Hu Z. The convenient synthesis of benzimidazole derivatives catalyzed by I<sub>2</sub> in aqueous media. *Journal of Heterocyclic Chemistry*. 2006;43: 773-775. DOI: 10.1002/jhet.5570430338
35. Bahrami K, Khodaei MM, Kaviani I. A simple and efficient one-pot synthesis of 2-substituted benzimidazoles. *Synthesis*. 2007; 547-550.
36. Nagawade RR, Shinde DB. BF<sub>3</sub>.OEt<sub>2</sub> promoted solvent free synthesis of benzimidazole derivatives. *Journal of the Chinese Chemical Society*. 2006; 17: 453-456. DOI: 10.1002/jccs.201100591.
37. Nagawade RR, Shinde DB. Zirconyl(IV) chloride-promoted synthesis of benzimidazole derivatives. *Russian Journal of Organic Chemistry*. 2006; 42: 453-454. DOI: 10.1134/S1070428006030201.

- [38] Mobinikhaledi A, Zendejdel M, Jamshidi FH. Zeolite-catalyzed synthesis of substituted benzimidazoles under solvent-free condition and microwave irradiation. *Synthesis and Reactivity in Inorganic, Metal-Organic and Nano-Metal Chemistry*. 2007; 37: 175-177. DOI: 10.1080/15533170701301496.
39. Varala R, Nasreen A, Enugala R, Adapa SR. L-proline catalyzed selective synthesis of 2-aryl-1-arylmethyl-1H-benzimidazoles. *Tetrahedron Letters*. 2007; 48: 69-72. DOI: 10.1016/j.tetlet.2006.11.010.
40. Rathod SB, Lande MK, Arbad BR. Synthesis, characterization and catalytic application of MoO<sub>3</sub>/CeO<sub>2</sub>-ZrO<sub>2</sub> solid heterogeneous catalyst for the synthesis of benzimidazole derivatives, *Bulletin Korean Chemical Society*. 2010; 31: 2835-2840. DOI: 10.5012/bkcs.2010.31.10.2835.
41. Gedye R, Smith F, Westaway K, Ali H, Baldisera L, Laberge L. The Use of Microwave-Ovens for rapid Organic-Synthesis. *Tetrahedron Letters*. 1986; 27: 279-282. DOI: 10.1016/S0040-4039(00)83996-9.
42. Bougrin K, Soufiaoui M. Nouvelle voie de synthèse des arylimidazole sous irradiation micro-ondes en milieu sec. *Tetrahedron Letters*. 1995; 36: 3683-3686. DOI: 10.1016/0040-4039(95)00611-F.
43. Bougrin K, Loupy A, Petit A, Daou B, Soufiaoui M. Nouvelle voie synthèse des 2-trifluorométhylarylimidazoles sur montmorillonite K10 en milieu sec sous micro-onde. *Tetrahedron*. 2001; 57: 163-168.
44. Bendale PM, Sun C-M. Rapid microwave-assisted liquid-phase combinatorial synthesis of 2-(arylamino)benzimidazoles. *J. Comb. Chem.* 2002; 4: 359-361. DOI: 10.1021/cc0200080.
45. Getvoldsen GS, Elander N, Stone-Elander AA. UV monitoring of microwave-heated reactions-A feasibility study. *Chemistry European Journal*. 2002; 8: 2255-2260. DOI: 10.1002/1522-3765(20020517)8:10<2255::AID-CHEM2255>3.0.CO;2-R.
46. Boufatah N, Gellis A, Maldonado J, Vanelle P. Efficient microwave-assisted synthesis of new sulfonylbenzimidazole-4,7-diones: heterocyclic quinones with potential antitumor activity. *Tetrahedron*. 2004; 60: 9131-9137. DOI: 10.1016/j.tet.2004.07.070.
47. Zhang W, Tempest P. Highly efficient microwave-assisted fluororous Ugi and post-condensation reactions for benzimidazoles and quinoxalinones. *Tetrahedron Letters*. 2004; 45: 6757-6760. DOI: 10.1016/j.tetlet.2004.07.039.
48. Getvoldsen G, Fredriksson A, Elander N, Stone-Elander S. Microwave-assisted cyclocondensation of 1,2-diaminobenzene with [4-<sup>18</sup>F]fluorobenzoic acid: microwave synthesis of 2-(4-<sup>18</sup>F)fluorophenyl)benzimidazole. *Journal of labelled compounds and radiopharmaceuticals*. 2004; 47: 139-145. DOI: 10.1002/jlcr.807.
49. Martinez-Palou R, Zepeda LG, Höpfl H, Montoya A, Guzmán-Lucero DJ, Guzmán J. Parallel and automated library synthesis of 2-long alkyl chain benzoazoles and azole[4,5-*b*]pyridines under microwave irradiation. *Molecular Diversity*. 2005; 9: 361-369. DOI: 10.1007/s11030-005-6357-5.
50. VanVilet DS, Gillespie P, Scicinski JJ. Rapid one-pot preparation of 2-substituted benzimidazoles from 2-nitroanilines using microwave conditions. *Tetrahedron Letters*. 2005; 46: 6741-6743. DOI: 10.1016/j.tetlet.2005.07.130.
51. Lin A-Y, Isome Y, Stewart E, Liu J-F, Yohannes D, Yu L. Microwave-assisted one step high-throughput synthesis of benzimidazoles. *Tetrahedron Letters*. 2006; 47: 2883-2886. DOI: 10.1016/j.tetlet.2006.02.127.
52. Jing X, Zhu Q, Xu F, Ren X, Li D, Yan C. Rapid one-pot preparation of 2-substituted benzimidazoles from esters using microwave conditions. *Synthetic Communication*. 2006; 36: 2601-2606. DOI: 10.1080/00397910600764204.
53. Gellis A, Kovacic H, Boufatah N, Vanelle. Synthesis and cytotoxicity evaluation of some benzimidazole-4,7-diones as bioreductive anticancer agents. *European Journal of Medicinal Chemistry*. 2008; 43: 1858-1864. doi:10.1016/j.ejmech.2007.11.020.

54. Algül O, Kaessler A, Apcin Y, Yilmaz A, Jose J. Comparative studies on conventional and microwave synthesis of some benzimidazole, benzothiazole and indole derivatives and testing on inhibition of hyaluronidase. *Molecules*. 2008; 13: 736-748.
55. Hosamani KM, Seetharamareddy HR, Keri RS, Hanamanthagouda MS, Moloney MG. Microwave assisted, one-pot synthesis of 5-nitro-2-aryl substituted-1H-benzimidazole libraries: Screening in vitro for antimicrobial activity. *Journal of Enzyme Inhibition and Medicinal Chemistry*. 2009; 24: 1095-1100. DOI: 10.1080/14756360802632716.
56. Hasaninejad A, Niknam K, Zare A, Farsimadan E, Shekouhy M. Silphox [POCl<sub>3-n</sub> (SiO<sub>2</sub>)<sub>n</sub>] as a new, efficient, and heterogeneous reagent for the synthesis of benzimidazole derivatives under microwave irradiation. *Phosphorus, Sulfur, and Silicon*. 2009; 184: 147-155. DOI: 10.1080/10426500802080931.
57. Chen C-H, Chien M-H, Kuo M-L, Chou C-T, Lai J-J, Lin S-F, Thummanagoti S, Sun C-M. Diversity-oriented synthesis of angular bis-benzimidazole derivatives under microwave irradiation. *Journal of Combinatorial Chemistry*. 2009; 11: 1038-1046. DOI: 10.1021/cc7002045.
58. Chen L-H, Hsiao Y-S, Yellol GS, Sun C-M. Microwave Promoted Simple, Efficient and Regioselective Synthesis of Trisubstituted Imidazo[1,2-*a*]benzimidazoles on Soluble Support. *ACS Combinatorial Science*. 2011 Mar 14;13(2):112-9. DOI:10.1021/co1000037.
59. Chanda K, Maiti B, Yellol GS, Chien M-H, Kuo M-L, Sun C-M. Polymer supported synthesis of novel benzoxazole linked benzimidazoles under microwave conditions: In vitro evaluation of VEGFR-3 kinase inhibition activity. *Organic & Biomolecular Chemistry*. 2011; 9: 1917-1926. DOI: 10.1039/c0ob00547a.
60. Chou C-T, Yellol GS, Chang W-J, Sun M-L, Sun C-M. Microwave assisted straightforward synthetic method for benzimidazole linked quinoxalinones on soluble polymer support. *Tetrahedron*. 2011; 67: 2110-2117. DOI: 10.1016/j.tet.2011.01.040.
61. Thummanagoti S, Yellol GS, Sun C-M. Ionic liquid supported multistep divergent synthesis of benzimidazole linked pyrrolo-/pyrido-/isoindolo-benzimidazolones. *Tetrahedron Letters*. 2011; 52: 2818-2822. DOI: 10.1016/j.tetlet.2011.03.046.
62. Ben-Alloum A, Bakkas S, Soufiaoui M. Benzimidazoles: Oxydation hétérocyclisante par le nitrobenzène ou le diméthylsulfoxyde sur silice et sous irradiation micro-ondes ou ultra-violet. *Tetrahedron Letters*. 1998; 39: 4481-4484.
63. Navarrete-Vazquez G, Moreno-Diaz H, Aguirre-Crespo F, Leon-Rivera I, Villalobos-Molina R, Munoz-Muniz O, Estrada-Soto S. Design, microwave-assisted synthesis, and spasmolytic activity of 2-(alkoxyaryl)-1H-benzimidazole derivatives as constrained stilbene bioisosteres. *Bioorganic & Medicinal Chemistry Letters*. 2006; 16: 4169-4173. DOI: 10.1016/j.bmcl.2006.05.082.
64. Zahran MA-H, El-Essawy FA-A, Yassin SM, Salem TA-R. Rapid and efficient synthesis of 4-substituted pyrazol-5-one under microwave irradiation in solvent-free conditions. *Archiv der Pharmazie*. 2007; 340: 591-598. DOI: 10.1002/ardp.200700121.
65. Dubey R, Moorthy NSHN. Comparative studies on conventional and microwave assisted synthesis of benzimidazole and their 2-substituted derivative with the effect of salt form of reactant. *Chemical Pharmaceutical Bulletin*. 2007; 55: 115-117. DOI: 10.1248/cpb.55.115.
66. Navarrete-Vazquez G, Moreno-Diaz H, Estrada-Soto S, Torres-Piedra M, Leon-Rivera I, Tlahuext H, Munoz-Muniz O, Torres-Gomez H. Microwave-assisted one-pot synthesis of 2-(substituted phenyl)-1H-benzimidazole derivatives. *Synthetic Communications*. 2007; 37: 2815-2825. DOI: 10.1080/003979107011473325.
67. Jacob RG, Dutra LG, Radatz CS, Mendes SR, Perin G, Lenardao EJ. Synthesis of 1,2-disubstituted benzimidazoles using SiO<sub>2</sub>/ZnCl<sub>2</sub>. *Tetrahedron Letters*. 2009; 50: 1495-1497. DOI: 10.1016/j.tetlet.2009.01.076.
68. Mao Z, Wang Z, Li J, Song X, Luo Y. Rapid and cheap synthesis of benzimidazoles via intermittent microwave promotion: A simple and potential industrial application of air as oxidant. *Synthetic Communication*. 2010; 40: 1963-1977. DOI: 10.1080/00397910903219328.

69. Azarifar D, Pirhayati M, Maleki B, Sanginabadi M, Yami RN. Acetic acid-promoted condensation of *o*-phenylenediamine with aldehydes into 2-aryl-1- (arylmethyl)-1H-benzimidazoles under microwave irradiation. *Journal of the Serbian Chemical Society*. 2010; 75: 1181-1189. DOI: 10.2298/JSC090901096A.
70. Mungra DC, Patel MP, Patel RG. Microwave-assisted synthesis of some new tetrazolo[1,5-*a*]quinoline-based benzimidazoles catalyzed by *p*-TsOH and investigation of their antimicrobial activity. *Medicinal Chemistry Research*. 2011; 20: 782-789. DOI: 10.1007/s00044-010-9388-0.
71. Krim J, Grünawal C, Taourirte M, Engels JW. Efficient microwave-assisted synthesis, antibacterial activity and high fluorescence of 5 benzimidazole-2'-deoxyuridines. *Biorganic and Medicinal Chemistry*. 2012; 20: 480-486. DOI: 10.1016/j.bmc.2011.10.041.
72. Reedy ACS, Rao PS, Venkataratnam RV. Fluoro organics: Facile synthesis of novel 2- or 4-trifluoromethyl-1H-arylo-1,5-diazepines, oxazepines, thiazepines, 2- (1,1,1-trifluoroacetyl)imidazoles, oxazoles and thiazoles. *Tetrahedron*. 1997; 53: 5847-5854. DOI: 10.1016/S0040-4020(97)00244-5.
73. Hulme C, Chappeta S, Griffith C, Lee Y-S, Dietrich J. An efficient solution Phase synthesis of triazadibenzoazulenones: 'designer isonitrile free' methodology enabled by microwaves. *Tetrahedron Letters*. 2009; 50: 1939-1942. DOI: 10.1016/j.tetlet.2009.02.099.
74. Wang S-L, Ding J, Jiang B, Gao Y, Tu S-J. Microwave-Assisted Solvent-Dependent Reaction: Chemoselective Synthesis of Quinoxalin-2(1*H*)-ones, Benzo[*d*]imidazoles and Dipeptides. *ACS Combinatorial Science*. 2011. Sep 12;13(5):572-7. DOI: 10.1021/co2001247.
75. Shaw AY, Medda F, Hulme C. Facile and rapid route for the synthesis of novel norstatine analogs via PADAM-cyclization methodology. *Tetrahedron Letters*. 2012; 53: 1313-1315. DOI: 10.1016/j.tetlet.2011.12.073.
76. Wen Xiaolan, El Bakali J, Deprez-Poulain R, Deprez B. Efficient propylphosphonic anhydride (®T3P) mediated synthesis of benzothiazoles, benzoxazoles and benzimidazoles. *Tetrahedron Letters*. 2012; 53: 2440-2443. DOI:10.1016/j.tetlet.2012.03.007.